

Central Limit Theorems for Star-Generators of the Infinite Symmetric Group

Claus Köstler

UCC - NUI

Probabilistic Operator Algebra Seminar

University California, Berkeley

1. May 2023

(Joint work with Jacob Campbell and Alexandru Nica)

- Revised Version of Slides -

Introduction & Motivation

Infinite symmetric group, star-generators, characters

General Notation

- $\mathbb{N} = \{1, 2, 3, \dots, n, \dots\}$, $[n] = \{1, 2, \dots, n\}$
- S_∞ is the **group of all finite permutations** on \mathbb{N} :

$$S_\infty = \left\{ \sigma: \mathbb{N} \rightarrow \mathbb{N} \left| \begin{array}{l} \sigma \text{ is bijective with } \sigma(n) \neq n \\ \text{for only finitely many } n \in \mathbb{N} \end{array} \right. \right\}$$

- The transpositions

$$\gamma_1 = (1, 2), \quad \gamma_2 = (1, 3), \quad \dots \quad \gamma_n = (1, n+1), \quad \dots$$

are called **star-generators** of S_∞ ; and $\gamma_0 = e$ denotes the identity.

- A **character** $\chi: S_\infty \rightarrow \mathbb{C}$ is a positive definite function which is constant on conjugacy classes and normalized, i.e. $\chi(e) = 1$.

A simple observation ...

Proposition ('exchangeability' of star-generators [Gohm & K '10])

Let χ be a character on S_∞ . Then one has

$$\chi(\gamma_{k_1} \gamma_{k_2} \cdots \gamma_{k_n}) = \chi(\gamma_{\sigma(k_1)} \gamma_{\sigma(k_2)} \cdots \gamma_{\sigma(k_n)})$$

for any $n \in \mathbb{N}$, $k_1, k_2, \dots, k_n \in \mathbb{N}$, and all $\sigma \in S_\infty$.

Proof.

$$\gamma_{\sigma(k)} = (1, \sigma(k) + 1) = (\tilde{\sigma}(1), \tilde{\sigma}(k + 1)) = \tilde{\sigma}(1, k + 1) \tilde{\sigma}^{-1}$$

for $\tilde{\sigma} \in S_\infty$ with $\tilde{\sigma}(1) = 1$ and $\tilde{\sigma}(k + 1) = \sigma(k) + 1$. □

- $(\mathbb{C}[S_\infty], \text{tr}_\chi)$ is a **tracial *-algebraic probability space**.
- The **noncommutative random variables** $(\gamma_n)_{n=1}^\infty$ are exchangeable.
- $(\gamma_n)_{n=1}^\infty$ **should be identically distributed** and '**conditional independent**' (in a certain sense in noncommutative probability).

A simple observation ...

Proposition ('exchangeability' of star-generators [Gohm & K '10])

Let χ be a character on S_∞ . Then one has

$$\chi(\gamma_{k_1} \gamma_{k_2} \cdots \gamma_{k_n}) = \chi(\gamma_{\sigma(k_1)} \gamma_{\sigma(k_2)} \cdots \gamma_{\sigma(k_n)})$$

for any $n \in \mathbb{N}$, $k_1, k_2, \dots, k_n \in \mathbb{N}$, and all $\sigma \in S_\infty$.

Proof.

$$\gamma_{\sigma(k)} = (1, \sigma(k) + 1) = (\tilde{\sigma}(1), \tilde{\sigma}(k + 1)) = \tilde{\sigma}(1, k + 1) \tilde{\sigma}^{-1}$$

for $\tilde{\sigma} \in S_\infty$ with $\tilde{\sigma}(1) = 1$ and $\tilde{\sigma}(k + 1) = \sigma(k) + 1$. □

- $(\mathbb{C}[S_\infty], \text{tr}_\chi)$ is a **tracial *-algebraic probability space**.
- The **noncommutative random variables** $(\gamma_n)_{n=1}^\infty$ are exchangeable.
- $(\gamma_n)_{n=1}^\infty$ is **identically distributed** and '**conditionally CS independent**' (see Gohm & K '10 for example).

Classical De Finetti Theorem

An infinite sequence of random variables $X \equiv (X_1, X_2, \dots)$ is

- **exchangeable** if

$$\mathbb{E}(X_{i_1} \cdots X_{i_n}) = \mathbb{E}(X_{\sigma(i_1)} \cdots X_{\sigma(i_n)})$$

for all $n \in \mathbb{N}$, $i_1, \dots, i_n \in \mathbb{N}$ and finite permutations σ on \mathbb{N} .



Theorem (De Finetti)

The following are equivalent:

- X is exchangeable*
- X is conditionally i.i.d.*

A Noncommutative De Finetti Theorem

Theorem (Gohm & K '09, K '10)

Let the pair (\mathcal{M}, φ) denote a von Neumann algebra \mathcal{M} equipped with a faithful normal (tracial) state φ . Then an **exchangeable sequence** $(x_n)_{n \geq 1} \subset \mathcal{M}$ is **identically distributed** and **CS-independent** over its **tail algebra**

$$\mathcal{T} := \bigcap_{n \geq 1} \text{vN}\{x_n, x_{n+1}, \dots\}$$

i.e.

$$E(xy) = E(x)E(y)$$

for any $x \in \text{vN}\{x_i \mid i \in I\}$ and $y \in \text{vN}\{x_i \mid i \in J\}$, where $I, J \subset \mathbb{N}$ are disjoint. Here E denotes the φ -preserving normal conditional expectation from \mathcal{M} onto \mathcal{T} .

Thoma's Theorem as a Quantum de Finetti Theorem

Theorem (Thoma '64, Kerov & Vershik '81, Okounkov '97)

An extreme character of S_∞ has the form:

$$\chi(\sigma) = \prod_{k=2}^{\infty} \left(\sum_{i=1}^{\infty} a_i^k + (-1)^{k-1} \sum_{j=1}^{\infty} b_j^k \right)^{m_k(\sigma)}$$

Here $m_k(\sigma)$ denotes the number of k -cycles in the permutation σ and the double sequence $(a_1, a_2, \dots; b_1, b_2, \dots)$ satisfy

$$a_1 \geq a_2 \geq \dots \geq 0, \quad b_1 \geq b_2 \geq \dots \geq 0, \quad \sum_{i=1}^{\infty} a_i + \sum_{j=1}^{\infty} b_j \leq 1.$$

- Gohm & K '10: **first operator algebraic proof!**
- **Key idea:** exchangeability of the sequence of star-generators

Algebraic CLTs

Theorem (von Waldenfels, Speicher, ...)

Let (\mathcal{A}, φ) be a $*$ -algebraic probability space and suppose that the sequence $(x_n = x_n^*)_{n=1}^\infty \subset \mathcal{A}$ satisfies:

- 1 exchangeability
- 2 singleton vanishing property, i.e. for all $n \in \mathbb{N}$,

$$\varphi(x_{\ell(1)}x_{\ell(2)} \cdots x_{\ell(n)}) = 0$$

for any $\ell: \{1, 2, \dots, n\} \rightarrow \mathbb{N}$ with $\#\ell^{-1}(\{m\}) = 1$ for some $m \in \mathbb{N}$.

Let $S_n = \frac{1}{\sqrt{n}}(x_1 + x_2 + \dots + x_n)$. Then one has

$$\lim_{n \rightarrow \infty} \varphi(S_n^k) = \sum_{\pi \in \mathcal{P}_2([k])} \varphi_\pi$$

where $\varphi_\pi = \varphi(x_{\ell(1)} \cdots x_{\ell(k)})$ with $\pi = \ker(\ell)$.

Central Limit Theorems for Star-Generators

- left regular character (Biane '95)
- block characters (K-Nica '21)
- beyond block characters (Campbell-K-Nica '22)

CLT for the character φ_{reg} of S_∞

$$(0, 0, \dots; 0, 0, \dots) \longleftrightarrow \varphi_{\text{reg}}(\sigma) := \begin{cases} 1 & \text{if } \sigma = e \\ 0 & \text{otherwise} \end{cases} \quad (\sigma \in S_\infty)$$

Theorem (Biane'95)

Let
$$s_n = \frac{1}{\sqrt{n}} (\gamma_1 + \gamma_2 + \dots + \gamma_n).$$

Then there exists $\mu_{\text{reg}} \in M_1(\mathbb{R})$ such that

$$\lim_{n \rightarrow \infty} \varphi_{\text{reg}}(s_n^k) = \int_{\mathbb{R}} t^k \mu_{\text{reg}}(dt) \quad (k \in \mathbb{N}).$$

Moreover, μ_{reg} equals Wigner's semicircle law.

CLT for a block character φ_d

$$\underbrace{\left(\frac{1}{d}, \frac{1}{d}, \dots, \frac{1}{d}, 0, 0, \dots; 0, 0, \dots\right)}_{d \text{ occurrences}} \longleftrightarrow \varphi_d(\sigma) := (1/d)^{\|\sigma\|} \quad (\sigma \in S_\infty)$$

Here $\|\sigma\| = \min\{m \mid \sigma = \tau_1 \tau_2 \cdots \tau_m \text{ for transpositions } \tau_1, \tau_2, \dots, \tau_m\}$

Theorem (K-Nica '21, Campbell-K-Nica '22)

Let
$$s_n = \frac{1}{\sqrt{n}} \left(\gamma_1 + \gamma_2 + \cdots + \gamma_n - \frac{n}{d} \right).$$

Then there exists a central limit law $\mu_d \in M_1(\mathbb{R})$ such that

$$\lim_{n \rightarrow \infty} \varphi_d(s_n^k) = \int_{\mathbb{R}} t^k \mu_d(dt) \quad (k \in \mathbb{N}).$$

Moreover, μ_d equals the average empirical distribution ν_d of a traceless random $d \times d$ GUE matrix with variance 1.

Traceless GUE Matrix

Consider a collection of d^2 independent centred Gaussian random variables

$$\{X_{i,j} \mid 1 \leq i \leq j \leq d\} \cup \{Y_{i,j} \mid 1 \leq i < j < d\}$$

with $\text{Var}(X_{ii}) = \frac{1}{d}$ for $1 \leq i \leq d$

and $\text{Var}(X_{ij}) = \text{Var}(Y_{ij}) = \frac{1}{2d}$ for $1 \leq i < j \leq d$.

Traceless Random $d \times d$ GUE Matrix of Variance 1

$$M = \begin{bmatrix} X_{11} & X_{12} + iY_{12} & \cdots & X_{1d} + iY_{1d} \\ X_{12} - iY_{12} & X_{22} & \cdots & X_{2d} + iY_{2d} \\ \vdots & \vdots & \ddots & \vdots \\ X_{1d} - iY_{1d} & X_{2d} - iY_{2d} & \cdots & X_{dd} \end{bmatrix} - \frac{X_{11} + \dots + X_{dd}}{d} I_d$$

with average empirical distribution ν_d (uniquely determined by

$\nu_d(M^k) = \mathbb{E}(\text{tr}_d(M^k))$ for all $k \in \mathbb{N}$).

The character $\varphi_{\underline{w}}$

For the Thoma double sequence $\underline{w} := (\underbrace{w_1, w_2, \dots, w_d}_{\text{summing up to 1}}, 0, 0, \dots; 0, 0, \dots)$,

$$\varphi_{\underline{w}}(\sigma) := \prod_{k=2}^{\infty} \left(\sum_{i=1}^d w_i^k \right)^{m_k(\sigma)} \quad (\sigma \in S_{\infty})$$

Postponing some notation

Theorem (Abstract CLT for Thoma character $\varphi_{\underline{w}}$)

Let $s_n = \frac{1}{\sqrt{n}} (U(\gamma_1) + U(\gamma_2) + \dots + U(\gamma_n) - nA_0)$.

Then there exists a central limit law $\mu_{\underline{w}} \in M_1(\mathbb{R})$ such that

$$\lim_{n \rightarrow \infty} \varphi_{\underline{w}}(s_n^k) = \int_{\mathbb{R}} t^k \mu_{\underline{w}}(dt) \quad (k \in \mathbb{N}).$$

... providing needed notation:

Let $U: S_\infty \rightarrow B(\mathcal{H})$ denote the GNS representation of $\varphi_{\underline{w}}$.

Proposition (Law of Large Numbers)

Let $\underline{w} = (w_1, w_2, \dots, w_d)$ be given. The operator

$$A_0 := \text{SOT} - \lim_{n \rightarrow \infty} \frac{1}{n} (U(\gamma_1) + \dots + U(\gamma_n)) \in B(\mathcal{H})$$

is a selfadjoint contraction with $\text{Spectrum}(A_0) = \bigcup_{i=1}^d \{w_i\}$.

Remark

- $A_0 = \text{WOT} - \lim_{n \rightarrow \infty} U(\gamma_n)$ ('limit-2-cycle $(1, \infty)$ ')
- $\mathcal{T} = \text{vN}\{A_0\}$ (tail algebra of the sequence $(U(\gamma_n))_{n=1}^\infty$)
- \mathcal{T} is trivial $\iff \text{Spectrum}(A_0) = \{1/d\}$

CCR-analogue of complex Gaussian random variable

Let (\mathcal{A}, φ) be a $*$ -probability space and $0 < \omega_{(1,*)}, \omega_{(*,1)} < \infty$ be fixed. An element $a \in \mathcal{A}$ is said to be a centred CCR-complex-Gaussian element if

$$a^*a = aa^* + (\omega_{(*,1)} - \omega_{(1,*)})1_{\mathcal{A}}$$

and, for $p, q \in \mathbb{N}_0$,

$$\varphi(a^p(a^*)^q) = \begin{cases} 0 & \text{if } p \neq q; \\ p! \omega_{(1,*)}^p & \text{if } p = q. \end{cases}$$

Remark: If $\omega_{(*,1)} = \omega_{(1,*)}$, then one recovers the usual notion of a complex Gaussian random variables.

Traceless CCR-GUE Matrix

A selfadjoint matrix $M = [a_{i,j}]_{1 \leq i,j \leq d} \in M_d(\mathcal{A})$ is called a traceless CCR-GUE matrix with parameters w_1, w_2, \dots, w_d if there exist commuting independent unital $*$ -subalgebras $\{\mathcal{A}_o\} \cup \{\mathcal{A}_{i,j} \mid 1 \leq i < j \leq d\} \subset \mathcal{A}$ s.t.:

- 1 For $1 \leq i < j \leq d$, the element $a_{i,j} \in \mathcal{A}_{i,j}$ is centred CCR-Gaussian (with parameters w_j and w_i) and $a_{j,i} := a_{i,j}^*$ such that

$$a_{j,i}a_{i,j} = a_{i,j}a_{j,i} + (w_j - w_i)1_{\mathcal{A}}$$

- 2 \mathcal{A}_o is commutative and the selfadjoint elements $a_{1,1}, \dots, a_{d,d} \in \mathcal{A}_o$ form a centred Gaussian family with covariance matrix

$$\begin{bmatrix} w_1 - w_1^2 & -w_1w_2 & \cdots & -w_1w_d \\ -w_1w_2 & w_2 - w_2^2 & \cdots & -w_2w_d \\ \vdots & \vdots & \ddots & \vdots \\ -w_1w_d & -w_2w_d & \cdots & w_d - w_d^2 \end{bmatrix}$$

Main Result

Theorem (Campbell-K-Nica '22)

Let (\mathcal{A}, φ) be a $*$ -probability space and let $M = [a_{i,j}]_{1 \leq i,j \leq d}$ be a traceless CCR-GUE matrix with parameters w_1, w_2, \dots, w_d . Consider the linear functional $\varphi_{\underline{w}}: M_d(\mathcal{A}) \rightarrow \mathbb{C}$ defined by

$$\varphi_{\underline{w}}(X) = \sum_{i=1}^d w_i \varphi(x_{i,i}) \quad \text{for } X = [x_{i,j}]_{i,j=1}^d \in M_d(\mathcal{A}).$$

Then the law of M in the $*$ -probability space $(M_d(\mathcal{A}), \varphi_{\underline{w}})$ is equal to the central limit law $\mu_{\underline{w}}$.

Rough Strategy for Proof of Main Result

- (1) Determine a moment formula for

$$\varphi_{\underline{w}}\left(\left(U(\gamma_{\ell(1)}) - A_0\right) \cdots \left(U(\gamma_{\ell(k)}) - A_0\right)\right)$$

for $\ell: [k] \rightarrow \mathbb{N}$ with $\ker(\ell) \in \mathcal{P}_2([k])$.







- (2a) Provide a Wick's lemma for a CCR-complex Gaussian element.
- (2b) Use this to get a formula for the joint moments of the entries of a CCR-GUE matrix.
- (3) Show that the formula from (1) and that from (2b) coincide.

For details see:

J. Campbell, C. Köstler, A. Nica; A central limit theorem for star-generators of S_∞ , which relates to traceless CCR-GUE matrices. *International Journal of Mathematics*, Volume 33, Issue 09, August 2022.

Thank you for your attention!

References

-  P. Biane. Permutation model for semi-circular systems and quantum random walks. *Pacific Journal of Mathematics* 171 (1995), 373-387.
-  J. Campbell, C. Köstler, A. Nica. A central limit theorem for star-generators of S_∞ , which relates to traceless CCR-GUE matrices. *Pacific Journal of Mathematics* 171 (1995), Paper No. 2250065.
-  R. Gohm, C. Köstler. Noncommutative independence from the braid group \mathbb{B}_∞ . *Communications in Mathematical Physics*, Volume 289, 435-482 (2009).
-  R. Gohm, C. Köstler. Noncommutative independence from characters of the infinite symmetric group S_∞ . Eprint arXiv:1005.5726 (2010).
-  C. Köstler. A noncommutative extended de Finetti theorem, *Journal of Functional Analysis* 258 (2010), 1073-1120.
-  C. Köstler, A. Nica. A central limit theorem for star-generators of S_∞ , which relates to the law of a GUE matrix, *Journal of Theoretical Probability* 34 (2021), 1248-1278.